

Vaxiduk

Freeze-dried pellet and solvent for suspension for injection

Ducks

Qualitative and quantitative composition

Each 0.5 ml-dose of vaccine contains:

- Live attenuated duck herpesvirus, Jansen strain $\geq 10^2$ CCID50^(*)
 - Aluminium hydroxide qs 0.85 mg
 - Excipient qs 0.5 ml
- ^(*) CCID50: 50 % cell culture infectious dose

Indication

In ducks, active immunisation against duck plague.

Immunological properties

The vaccine contains a strain of duckling herpesvirus virus and is intended to provide an active immunisation against duck plague in healthy and contaminated environment.

Dosage for each species, route(s) and method of administration

Inject one 0.5-ml dose per animal, by subcutaneous or intramuscular route, according to the following schedule:

Primary vaccination

2 injections with a 4-week interval between the 7th and 11th week of age.

Boosters

1 injection prior to each laying season.

In contaminated environment, birds of at least 10 days of age should be administered one dose of vaccine as from the first symptoms.

Apply usual aseptic procedures. Use antiseptic- or disinfectant-free material for injection purposes. Shake the reconstituted vaccinal suspension before use.

Contra-indications

- Do not vaccinate breeders during moulting.
- Do not administrate simultaneously with another live virus vaccine.

Special warning for each target species

In ducklings, the presence of maternally derived antibodies may interfere with the post-vaccination immune response. The vaccine strain may spread.

Special precautions to be taken by the person administering the medicinal product to animals

If the vaccine is accidentally injected to man, urgent medical attention is necessary.

Use during pregnancy, lactation or lay

Do not vaccinate birds in lay.

Interactions with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be decided on a case by case basis.

Overdose (symptoms, emergency procedures, antidotes)

No undesirable effect was observed following the administration of 10 doses of vaccine.

Withdrawal period: Zero days.

Special precautions for storage

Freeze-dried pellet: store and transport at a temperature between +2°C and +8°C, protected from light.

Solvent: do not store at a temperature above 25°C. Do not freeze.

After reconstitution, use immediately.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

For animal treatment only.



MERIAL GRAPHIC ARTWORK APPROVAL

Code: 0 053 914425	VERSION	PTG : 0041 Version : B 040516	COLORS
PRODUCT: VAXIDUK NOT VERSO	INITIALS: VA - PICTURAL AN		Noir/Black
COUNTRY: FRANCE/EXPORT (FR/GB)	VERSION - DATE: A 20/09/16 - 9h35		
Dimensions à plat (mm): 175 x 114	H:020316		
Dimensions plié(s): 20 x 114			
PRINTED BASIS CODE (if applicable): NA	CANCEL AND REPLACE: 0 053 902419		! Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:	Regulatory approval:
COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:
	veroniq ue.abatt u2@san ofi.com Signature numérique de veronique.abattu2@sanofi.com DN: cn=veronique.aba ttu2@sanofi.com Date: 2016.10.05

NOT PRINTABLE
NON IMPRIMABLE

TECHNICAL NOTE(S) - INFO(S) TECHNIQUE

- Artwork identification(s)
- Text areas / zones de textes
- Dieline / Découpe