


# Surveillance of Influenza A virus in Danish pigs

## Samples and results 2023

The table illustrate the number of samples, submissions and herds that contributed to the surveillance program each month. In addition, the results of the influenza A virus and H1pdm09 screening are shown.


	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<b>Received</b>												
Samples	169	118	163	143	89	101	97	136	144			
Submissions	50	34	49	46	27	31	27	42	42			
Herds	46	34	48	45	27	31	27	41	42			
<b>Influenza positive</b>												
Samples	84	79	67	61	54	45	44	52	76			
Submissions	32	26	27	25	20	18	18	26	25			
Herds	31	26	27	25	20	18	18	26	25			
<b>H1pdm09 positive</b>												
Samples	36	32	25	12	31	21	15	16	24			
Submissions	15	11	13	6	12	8	7	12	12			
Herds	15	11	13	6	12	8	7	12	12			

In September, 42 submissions from 42 herds registered with different CHR numbers were received. On average, one submission included 3.4 samples. The positive percentage for the month of September was 59 %, which is comparable to June and August. All influenza A virus positive samples were tested for the presence of H1pdm09. In total, 48 % of the influenza A virus positive submissions were positive for H1pdm09 resembling the proportion observed previously.



The figure illustrates the percentage of influenza A virus negative and positive submissions including the proportion of H1pdm09 positive submissions.


## Distribution of swine influenza A virus subtypes



Swine influenza A virus can be classified into subtypes and genotypes. The subtype describes the combination of HA and NA surface gene segments, and the genotype describes the combination of all eight segments based on their genetic origin. Information on the contemporary circulating swine influenza A virus subtypes is essential for the update of vaccination protocols, optimization of the diagnostic assays and for evaluation of the zoonotic risk.

In September, the subtype (HA and NA gene segments) was determined for 21 submissions. The H1avNx viruses were dominating, with H1avN2sw being the most prevalent. In addition, a major proportion of the submissions included H1pdm09N1av followed by other H1pdm09N1x viruses, similar to the pattern observed in August and March.

## Distribution of swine influenza A virus genotypes



For September, 18 submissions were genotyped. For the H1avN2sw viruses, the dominating genotype was H1avN2sw-1 which contained all gene segments of avian-like origin, followed by H1avN2sw-3. For the H1pdmNx viruses (H1pdmN1av, H1pdmN1pdm, H1pdmN2sw), six samples were genotyped with H1pdmN1av-2 which had an internal gene cassette of H1N1pdm09 origin except from the NS segment that was of avian-like H1N1 origin.

# Phylogenetic analysis

